

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

VERBENA PLANT NAMED 'SUNMARISAKURA'**BOTANICAL CLASSIFICATION***Verbena hybrida***VARIETAL DENOMINATION**

5 'Sunmarisakura'

BACKGROUND OF THE VARIETY

10 The present new distinct cultivar of *Verbena* was created by heavy ion beam irradiation of the *Verbena* hybrid variety called 'Sunmarisa'. 'Sunmarisa' (U.S. Plant Patent No. 11130) is our *Verbena* hybrid variety grown at Yokaichi-shi, Shiga-ken, Japan.

15 In July 1998, 5Gy of ionic nitrogen (135MeV) was irradiated onto 128 pieces of *In vitro* axillary bud of 'Sunmarisa' using the Ring Cyclotron at The Institute of Physical and Chemical Research. Two weeks later, elongated buds were grown from cuttings. In March 1999, five varieties were selected in view of self-sterility. These plants were propagated by cutting and then grown in pot and planter box on trial from May 1999 at Yokaichi-shi, Shiga-ken, Japan. Additionally, the plants were grown in pots and planter boxes during trial in 2000. Finally, one plant was selected from these five varieties in view of having a long flower duration. By November 2000, the botanical characteristics of the selected plant were examined. As a result, it was concluded that this new *Verbena* plant is distinguishable from any other variety, whose existence is known to us, and is uniform and stable in its characteristics. This new variety of *Verbena* plant was named 'Sunmarisakura'.

25 The new variety of the present invention can be distinguished from its parent 'Sunmarisa' in its self-sterility. Also the new variety can be distinguished from the similar variety 'Sunmaripi' in the flower color and self-sterility.

The new variety of *Verbena* plant 'Sunmarisakura' was asexually reproduced by cutting at Yokaichi-shi, Shiga-ken, Japan, and the homogeneity and stability

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

thereof were confirmed. The instant plant retains its distinctive characteristics and reproduces true to type in successive generations.

In the following description, the color-coding is in accordance with The Horticultural Colour Chart of The Royal Horticultural Society, London, England (R.H.S.).

'Sunmarisa' was used to produce the new variety, 'Sunmarisakura'. The botanical characteristics of 'Sunmarisa' are as follows when grown at Yokaichi-shi, Shiga-ken, Japan.

Plant:

- 10 Growth habit.—Decumbent.
 Plant width.—Approximately 65-71 cm.
 Plant height.—Approximately 8-9 cm.

Stem:

- Diameter.—Approximately 1.7-2.2 mm.
15 Anthocyanin pigmentation.—Absent.
 Pubescence.—Normal.
 Branching.—A moderate level of branching is present.
 Subterranean stem.—Absent but when the stems contact the surface of soil, the nodes take root in the ground and the plant growth thereby spreads.
20 Length of internodes—Approximately 1.7-2.7 cm.

Leaf:

- Phyllotaxis.—Opposite.
 Shape.—Oblong-lanceolate.
 Depth of incision.—Shallow.
25 Margin.—Serrate.
 Length.—Approximately 2.9-3.6 cm.
 Width.—Approximately 1.6-2.4 cm.
 Leaf apex.—Mucronate.

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

Leaf base.—Petiolate.

Color.—Dark olive green (near R.H.S. 141A) on the upper surface.

Pubescence.—Sparse.

Petiole.—Present.

5 Thickness of petiole.—Approximately 1.0-1.2 mm.

Length of petiole.—Approximately 1.2-2.5 mm.

Flower:

Shape of cluster.—Obovate.

Length of cluster.—Approximately 2.3-4.4 cm.

10 Diameter of cluster.—Approximately 4.0-5.8 cm.

Facing direction.—Upward.

Outward curvature of petal.—Flat.

Diameter.—Approximately 1.8-1.9 cm.

Height.—Approximately 2.1-2.2 cm.

15 Color.—Light purplish pink (near R.H.S. 62C) on the upper surface.

Eye color.—Pale yellow (near R.H.S. 11D).

Variegation on petal.—Absent.

Color presentation.—Substantially even.

Color intensity.—Absent.

20 Overlapping of petals.—Separate.

Incision of petal.—Present.

Number of petals.—5.

Sepal.—Tubular in configuration.

Length of calyx.—Approximately 1.2-1.4 cm.

25 Anthocyanin pigmentation of calyx limb.—Absent.

Shape of pistil.—Two lobes.

Color of anther.—Yellowish green.

Diameter of peduncle.—Medium.

Length of peduncle.—Medium.

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

Number of flowers.--Many.

Flower bearing.--In a cluster.

Reproductive organs.--1 pistil and 4 stamens.

Pollen.--Brilliant greenish-yellow (near R.H.S. 6C) in coloration.

5 Flower fragrance.--Absent.

Flowering time.--Early.

Flowering duration.--Long. When planted during March, the plant commonly blooms from April to November. A bloom cluster commonly is present for approximately 2 to 3 weeks, and an individual bloom within the cluster commonly lasts for approximately 7 to 10 days on the plant. The plant is highly tolerant to heat, and exhibits a high resistance to diseases, particularly powdery mildew, and a high resistance to pests.

'Sunmaripi' was used as a comparison for this new variety 'Sunmarisakura'. The botanical characteristics of the 'Sunmaripi' are as follows when grown at when grown at Yokaichi-shi, Shiga-ken, Japan.

Plant:

Growth habit.--Decumbent.

Plant width.--Approximately 60 cm.

Plant height.--Approximately 20 cm.

Stem:

Diameter.--Approximately 2.5 mm.

Anthocyanin pigmentation.--Present.

Pubescence.--Normal.

Branching.--A moderate level of branching is present.

Subterranean stem.--Absent but when the stems contact the surface of soil, the nodes take root in the ground and the plant growth thereby spreads.

Length of internode.--Approximately 4.5 cm.

Leaf:

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

Phyllotaxis.—Opposite.

Shape.—Oblong-lanceolate.

Depth of incision.—Shallow.

Margin.—Serrate.

5 Length.—Approximately 3.5 cm.

Width.—Approximately 2.2 cm.

Leaf apex.—Mucronate.

Leaf base.—Petiolate.

Color.—Grayish olive green (near R.H.S. 147A) on the upper surface.

10 Pubescence.—Sparse.

Petiole.—Present.

Thickness of petiole.—Approximately 1.0 mm.

Length of petiole.—Approximately 2.0 mm.

Flower:

15 Shape of cluster.—Obovate.

Length of cluster.—Approximately 3.2 cm.

Diameter of cluster.—Approximately 5.2 cm.

Facing direction.—Upward.

Outward curvature of petal.—Flat.

20 Diameter.—Approximately 1.7 cm.

Height.—Approximately 2.0 cm.

Color.—Deep purplish pink (near R.H.S. 77B) on the upper surface.

Eye color.—Pale yellow (near R.H.S. 11D).

Variegation on petal.—Absent.

25 Color presentation.—Substantially even.

Color intensity.—Absent.

Incision of petal.—Present.

Number of petals.—5.

Sepal.—Tubular in configuration.

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

Length of calyx.—Approximately 1.0 cm.

Anthocyanin pigmentation of calyx limb.—Present.

Shape of pistil.—Two lobes.

Color of anther.—Yellowish green.

5 Diameter of peduncle.—Medium.

Length of peduncle.—Medium.

Number of flowers.—Many.

Flower bearing.—In a cluster.

Reproductive organs.—1 pistil and 4 stamens.

10 Pollen.—Brilliant greenish-yellow (near R.H.S. 6C) in coloration.

Flower fragrance.—Absent.

Flowering time.—Early.

Flowering duration.—Long. When planted during March, the plant commonly blooms from April to November. A bloom cluster commonly is present for approximately 2 to 3 weeks, and an individual bloom within the cluster commonly lasts for approximately 7 to 10 days on the plant. The plant is highly tolerant to heat, and exhibits a high resistance to diseases, particularly powdery mildew, and a high resistance to pests.

15

SUMMARY OF THE NEW VARIETY

20

The new variety of *Verbena* plant has a broad spreading growth habit and long stems. The plant is well branched and abundantly forms flowers in a cluster. The blooms are present in profusion. The blooming period is April to November and flowering duration is long. The entire plant remains in bloom for an extended period of time. The flower size is large and the petal coloration of the flowers is light purplish pink. The plant is highly tolerant to heat, exhibits a high resistance to pests and diseases, particularly powdery mildew. The plant has very low self-fertility

25

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts a side view of the 'Sunmarisakura' plant of the present invention.

FIG. 2 depicts a close-up view of the flowers of the 'Sunmarisakura' plant of the present invention.

DESCRIPTION OF THE VARIETY

The botanical characteristics of the new and distinct variety of *Verbena* plant, 'Sunmarisakura' are as follows when grown at Yokaichi-shi, Shiga-ken, Japan.

Plant:

Growth habit.—Decumbent.

Plant width.—Approximately 65-71 cm.

Plant height.—Approximately 8-9 cm.

Stem:

Diameter.—Approximately 1.7-2.2 mm.

Anthocyanin pigmentation.—Absent.

Pubescence.—Normal.

Branching.—A moderate level of branching is present.

Subterranean stem.—Absent but when the stems contact the surface of soil, the nodes take root in the ground and the plant growth thereby spreads.

Length of internode.—Approximately 1.7-2.7 cm.

Leaf:

Phyllotaxis.—Opposite.

Shape.—Oblong-lanceolate.

Depth of incision.—Shallow.

Margin.—Serrate.

Length.—Approximately 2.9-3.6 cm.

Width.—Approximately 1.6-2.4 cm.

Leaf apex.—Mucronate.

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

Leaf base.—Petiolate.

Color.—Dark olive green (near R.H.S. 141A) on the upper surface. Near
R.H.S. 138A on the lower surface.

Pubescence.—Sparse.

5 Petiole.—Present.

Thickness of petiole.—Approximately 1.0-1.2 mm.

Length of petiole.—Approximately 1.2-2.5 mm.

Flower:

Shape of cluster.—Obovate.

10 Length of cluster.—Approximately 2.3-4.4 cm.

Diameter of cluster.—Approximately 4.0-5.8 cm.

Facing direction.—Upward.

Outward curvature of petal.—Flat.

Diameter.—Approximately 1.8-1.9 cm.

15 Height.—Approximately 2.1-2.2 cm.

Color.—Light purplish pink (near R.H.S. 62C) on the upper surface. Near
R.H.S. 65A on the lower surface.

Eye color.—Pale yellow (near R.H.S. 11D).

Eye size.—Small.

20 Variegation on petal.—Absent.

Color presentation.—Substantially even.

Color intensity.—Absent.

Incision of petal.—Present.

Number of petals.—5.

25 Sepal.—Tubular in configuration.

Length of calyx.—Approximately 1.2-1.4 cm.

Anthocyanin pigmentation of calyx limb.—Absent.

Shape of pistil.—Two lobes.

Color of anther.—Yellowish green (near R.H.S. N144D).

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

Peduncle length.—Approximately 2.1 cm.

Peduncle diameter.—Approximately 1.5 mm.

Flower bearing.—In a cluster.

Reproductive organs.—1 pistil and 4 stamens.

5 Pollen.—Scarce. Brilliant greenish-yellow (Near R.H.S. 6C) in coloration.

Stigma color. — Near R.H.S. 144A

Style color. — Near R.H.S. 144C

Ovary color. — Near R.H.S. 144B

Flower fragrance.—Absent.

10 Flowering time.—Early.

Flowering duration.—Long. When planted during March, the plant commonly blooms from April to November. A bloom cluster commonly is present for approximately 2 to 3 weeks, and an individual bloom within the cluster commonly lasts for approximately 7 to 10 days on the plant. The plant is highly tolerant to
15 heat. The plant grows well at temperatures up to at least 35°C, and exhibits a high resistance to diseases, particularly powdery mildew. No serious damage by pathogens and pests common to *Verbena* has been observed.

This new variety of *Verbena* plant is most suitable for flower bedding and potting, particularly in planters, and is further excellent for use in a hanging basket.

20 The plant of this new variety, 'Sunmarisakura', is presently planted and maintained at Yokaichi-shi, Shiga-ken, Japan.

**SUBSTITUTE SPECIFICATION
(CLEAN COPY)**

52011/CAB/A400

It is claimed:

1. A new and distinct variety of *Verbena hybrida* plant named 'Sunmarisakura', substantially as herein illustrated and described.